## Computational approach for detecting undetected interstellar molecules: CH<sub>2</sub>DOD and CHD<sub>2</sub>OD

A. Badawi

Zewail City of Science and Technology, Egypt

Observations revealed that complex molecules (COMs) are very abundant in star forming regions (SFRs) either low-mass or massive. Methanol is believed to be the main driver of molecular complexity in the interstellar medium. Some deuterated forms of methanol; namely CH<sub>2</sub>DOD and CHD<sub>2</sub>OD, could play a role in the formation of deuterated COMs. We found that these undetected forms are detectable in SFRs (with abundances  $>10^{-12}n_H$ ). We ran theoretical models and computed the peak positions of the IR spectral lines of these species and compared them with experimentally predicted IR positions.

Our results are in agreement with experimental data. Given the calculated abundances of these undetected species and their spectral line positions, we recommend searches for these molecules in SFRs.

## Z. Awad<sup>1</sup> and <u>A. Badawi<sup>2</sup></u>

## <sup>1</sup>Astronomy, Space Science and Meteorology Department, faculty of science, Cairo University, Egypt

<sup>2</sup>Center for Photonics and Smart Materials, Zewail City of Science and Technology, Egypt

Observations revealed that complex molecules (COMs) are very abundant in star forming regions (SFRs) either low-mass or massive. Methanol is believed to be the main driver of molecular complexity in the interstellar medium. Some deuterated forms of methanol; namely CH<sub>2</sub>DOD and CHD<sub>2</sub>OD, could play a role in the formation of deuterated COMs. We found that these undetected forms are detectable in SFRs (with abundances  $>10^{-12}n_H$ ). We ran theoretical models and computed the peak positions of the IR spectral lines of these species and compared them with experimentally predicted IR positions.

Our results are in agreement with experimental data. Given the calculated abundances of these undetected species and their spectral line positions, we recommend searches for these molecules in SFRs.